

X-ray Thomson Scattering from Dense Plasmas

S. Glenzer

May 18, 2007

49the Annual Meeting of the Division of Plasma Physics orlando, FL, United States November 12, 2007 through November 16, 2007

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Tutorial talk nomination for the 49th Annual Meeting of the Division of Plasma Physics, November 12-16, 2007 , Rosen Centre Hotel , Orlando, Florida

X-ray Thomson scattering in dense plasmas*

O. L. Landen

Lawrence Livermore National Laboratory, USA

Advances in the development of laser-produced x-ray sources have enabled a new class of high-energy density physics experiments. Powerful narrow-bandwidth x rays penetrate through short-lived hot dense states of matter and probe the physical properties with spectrally resolved x-ray scattering. Experiments from isochorically-heated plasmas with electron densities in the range of solid density and above have been demonstrated allowing for the first time exploration of the microscopic properties of dense matter regime close to strongly-coupled and Fermi degenerate conditions. measurements have accessed the non-collective Compton scattering regime, which provides accurate diagnostic information on the temperature, density and ionization states. The forward scattering spectrum has been shown to measure the collective plasmon oscillations. Besides extracting the standard plasma parameters, density and temperature, forward scattering yields new observables such as a direct measure of collisions, quantum effects and detailed balance. In this talk, we will discuss new results important for applications of this technique for novel experiments in a wide range of research areas such as inertial confinement fusion, radiation-hydrodynamics, material science, and laboratory astrophysics.

This work was performed under the auspices of the U.S. Department of Energy by University of California Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

Supporting remarks:

Dr. Landen is the Associate Program Leader for Target Physics in the National Ignition Facility Program at the Lawrence Livermore National Laboratory. Dr Landen is a fellow of the American Physical Society and has led numerous laser experiments on the development and applications of x-ray sources. Dr. Landen first proposed x-ray Thomson scattering for probing Warm Dense Matter and has overseen the implementation of this novel experimental method at the Omega Laser Facility as senior scientist.

In this tutorial talk, Dr. Landen will present the extension of scattering theory and experimental techniques from traditional optical Thomson Scattering to the x-ray regime, including providing a historical perspective from the early days of Compton scattering off room temperature solids. He will relate the experimental observables to dynamic structure factor calculations and discuss the results in terms of various physics models.

Deleted:

He will further provide an outlook discussing future applications, e.g., for probing extremely dense states of matter in future experiments on the National Ignition Facility.

Publications

- O. L. Landen et al., Rev. Sci. Instrum. 72, 627 (2001).
- O. L. Landen, S.H. Glenzer, M.J. Edwards et al., JOSRT 71, 465 (2001).
- S. H. Glenzer, G. Gregori, O. L. Landen et al., Phys. Rev. Letters 90, 175002 (2003).
- G. Gregori, S. H. Glenzer, and O. L. Landen, J. Phys. A 36, 5971 (2003).
- G. Gregori, S. H. Glenzer and O. L. Landen, Phys. Rev. E 74, 026402 (2006).
- S. H. Glenzer, O. L. Landen, P. Neumayer et al., Phys. Rev. Lett. 98, 065002 (2007).

Informed individuals:

G, Gregori, CLF - Rutherford Appleton Laboratory, Chilton, Didcot, OX11 0QX Great Britain. Tel: +44 (0)1235 44 5215. "Gregori, G \(Gianluca\)" G.Gregori@rl.ac.uk

M. Koenig, Laboratoire LULI, Ecole Polytechnique, 91128 Palaiseau Cedex, FRANCE. Tel: 33-(0)169335399. Michel KOENIG michel.koenig@polytechnique.fr